Esperion Announces Submission of Supplemental New Drug Applications to U.S. FDA for NEXLETOL® (bempedoic acid) Tablet and NEXLIZET® (bempedoic acid and ezetimibe) Tablet

June 1, 2023

– Submitted four efficacy supplements seeking inclusion of CV risk reduction and expansion of LDL-C lowering –

– Anticipated approvals in U.S. in the first half of 2024 –

ANN ARBOR, Mich., June 01, 2023 (GLOBE NEWSWIRE) -- Esperion (NASDAQ: ESPR) today announced that it has submitted Supplemental New Drug Applications (sNDA) to the U.S. Food and Drug Administration (FDA) for the Company's oral non-statin products NEXLETOL and NEXLIZET. In its sNDAs, Esperion seeks to add the use of both NEXLETOL and NEXLIZET for cardiovascular risk reduction and also seeks to remove the statin limitation in the LDL-C indication. The sNDA submissions are based on the landmark Cholesterol Lowering via Bempedoic acid, an ACL-Inhibiting Regimen (CLEAR) Outcomes trial which showed NEXLETOL demonstrated significant cardiovascular risk reduction across a range of primary and second endpoints, including a 27% risk reduction of non-fatal myocardial infarction, a 23% risk reduction of the composite of fatal and non-fatal myocardial infarction, a 19% risk reduction of coronary revascularization, a 15% risk reduction of the MACE-3 composite, and a 13% risk reduction of the MACE-4 composite. Following the FDA's review of the submission, it will notify the Company if the applications are deemed complete for review by mid-August 2023. The Company anticipates FDA approval of the sNDAs in the first half of 2024.

"These submissions mark a pivotal achievement for Esperion as we seek to meaningfully expand the eligible patient populations for NEXLETOL and NEXLIZET," said Sheldon Koenig, President and Chief Executive Officer of Esperion. "Since the announcement of the CLEAR Outcomes results, we've seen accelerating adoption of our practice-changing treatments as prescribers, patients and payers alike recognize NEXLETOL and NEXLIZET as the clear next step after statins, and value their potential to significantly reduce cardiovascular risk."

INDICATION
NEXLETOL and NEXLIZET are indicated as adjuncts to diet and maximally tolerated statin therapy for the treatment of adults with heterozygous familial hypercholesterolemia or established atherosclerotic cardiovascular disease who require additional lowering of LDL-C. *Limitations of Use*: The effect of NEXLETOL and NEXLIZET on cardiovascular morbidity and mortality has not been determined.

IMPORTANT SAFETY INFORMATION
Contraindications: NEXLETOL has no contraindications. NEXLIZET is contraindicated in patients with a known hypersensitivity to ezetimibe tablets. Hypersensitivity reactions including anaphylaxis, angioedema, rash, and urticaria have been reported with ezetimibe.

Warnings and Precautions: Hyperuricemia: Bempedoic acid, a component of NEXLETOL and NEXLIZET, may increase blood uric acid levels. Hyperuricemia may occur early in treatment and persist throughout treatment, and may lead to the development of gout, especially in patients with a history of gout. Assess uric acid levels periodically as clinically indicated. Monitor for signs and symptoms of hyperuricemia, and initiate treatment with urate-lowering drugs as appropriate.
Tendon Rupture: Bempedoic acid is associated with an increased risk of tendon rupture or injury. In clinical trials, tendon rupture occurred in 0.5% of patients treated with bempedoic acid versus 0% of patients treated with placebo, and involved the rotator cuff (the shoulder), biceps tendon, or Achilles tendon. Tendon rupture occurred within weeks to months of starting bempedoic acid. Tendon rupture may occur more frequently in patients over 60 years of age, patients taking corticosteroid or fluoroquinolone drugs, patients with renal failure, and patients with previous tendon disorders. Discontinue NEXLETOL or NEXLIZET at the first sign of tendon rupture. Avoid NEXLETOL and NEXLIZET in patients who have a history of tendon disorders or tendon rupture.

Adverse Reactions: In NEXLETOL clinical trials, the most commonly reported adverse reactions were upper respiratory tract infection, muscle spasms, hyperuricemia, back pain, abdominal pain or discomfort, bronchitis, pain in extremity, anemia, and elevated liver enzymes. Reactions reported less frequently, but still more often than with placebo, included benign prostatic hyperplasia and atrial fibrillation.
In the NEXLIZET clinical trial, the most commonly reported adverse reactions observed with NEXLIZET, but not observed in clinical trials of bempedoic acid or ezetimibe, a component of NEXLIZET, and occurring more frequently than with placebo, were urinary tract infection, nasopharyngitis, and constipation.
Adverse reactions reported in clinical trials of ezetimibe, and occurring at an incidence greater than with placebo, included upper respiratory tract infection, diarrhea, arthralgia, sinusitis, pain in extremity, fatigue, and influenza. Other adverse reactions reported in postmarketing use of ezetimibe included hypersensitivity reactions, including anaphylaxis, angioedema, rash, and urticaria; erythema multiforme; myalgia; elevated creatine phosphokinase; myopathy/rhabdomyolysis; elevations in liver transaminases; hepatitis; abdominal pain; thrombocytopenia; pancreatitis; nausea; dizziness; paresthesia; depression; headache; cholelithiasis; cholecystitis.

Drug Interactions: Simvastatin and Pravastatin: Concomitant use with bempedoic acid results in increased concentrations and increased risk of simvastatin or pravastatin-related myopathy. Use of either NEXLETOL or NEXLIZET with greater than 20 mg of simvastatin or 40 mg of pravastatin should be avoided.
Cyclosporine: Caution should be exercised when using NEXLIZET and cyclosporine concomitantly due to increased exposure to both ezetimibe and cyclosporine. Monitor cyclosporine concentrations in patients receiving NEXLIZET and cyclosporine. In patients treated with cyclosporine, the potential effects of the increased exposure to ezetimibe from concomitant use should be carefully weighed against the benefits of alterations in lipid levels provided by NEXLIZET.

Fibrates: Coadministration of NEXLIZET with fibrates other than fenofibrate is not recommended. Fenofibrate and ezetimibe may increase cholesterol excretion into the bile, leading to cholelithiasis. If cholelithiasis is suspected in a patient receiving NEXLIZET and fenofibrate, gallbladder studies are indicated and alternative lipid-lowering therapy should be considered.
Cholestyramine: Concomitant use of NEXLIZET and cholestyramine decreases ezetimibe concentration. This may result in a reduction of efficacy. Administer NEXLIZET either at least 2 hours before, or at least 4 hours after, bile acid sequestrants.

Lactation and Pregnancy: It is not recommended that NEXLETOL or NEXLIZET be taken during breastfeeding. Discontinue NEXLETOL or NEXLIZET when pregnancy is recognized, unless the benefits of therapy outweigh the potential risks to the fetus. Based on the mechanism of action of bempedoic acid, NEXLETOL and NEXLIZET may cause fetal harm.

Please see full Prescribing Information here.

CLEAR Cardiovascular Outcomes Trial
CLEAR Outcomes is part of the CLEAR clinical research program for NEXLETOL® (bempedoic acid) Tablet and NEXLIZET® (bempedoic acid and ezetimibe) Tablet. The CLEAR Program seeks to generate important clinical evidence on the safety and efficacy of bempedoic acid, a first in a class ATP citrate lyase inhibitor contained in NEXLETOL and NEXLIZET and its potential role in addressing additional critical unmet medical needs. More than 60,000 people will have participated in the program by the time of its completion. The CLEAR Program includes 5 label-enabling Phase III studies as well as other key Phase IV studies with the potential to reach more than 70 million people with or at risk for CVD based on elevated LDL-C.

Esperion Therapeutics
At Esperion, we discover, develop, and commercialize innovative medicines to help improve outcomes for patients with or at risk for cardiovascular and cardiometabolic diseases. The status quo is not meeting the health needs of millions of people with high cholesterol – that is why our team of passionate industry leaders is breaking through the barriers that prevent patients from reaching their goals. Providers are moving toward reducing LDL-cholesterol levels as low as possible, as soon as possible; we provide the next steps to help get patients there. Because when it comes to high cholesterol, getting to goal is not optional. It is our life's work. For more information, visit esperion.com and esperionscience.com and follow us on Twitter at twitter.com/EsperionInc.

Forward-Looking Statements
This press release contains forward-looking statements that are made pursuant to the safe harbor provisions of the federal securities laws, including statements regarding regulatory submissions and potential approvals, marketing strategy and commercialization plans, current and planned operational expenses, future operations, commercial products, clinical development, including the timing, designs and plans for the CLEAR Outcomes study and its results, plans for potential future product candidates, financial condition and outlook, including expected cash runway, and other statements containing the words "anticipate," "believe," "estimate," "expect," "intend," "may," "plan," "predict," "project," "suggest," "target," "potential," "will," "would," "could," "should," "continue," and similar expressions. Any express or implied statements contained in this press release that are not statements of historical fact may be deemed to be forward-looking statements. Forward-looking statements involve risks and uncertainties that could cause Esperion's actual results to differ significantly from those projected, including, without limitation, the impact of the ongoing COVID-19 pandemic on our business, revenues, results of operations and financial condition, the net sales, profitability, and growth of Esperion's commercial products, clinical activities and results, supply chain, commercial development and launch plans, the outcomes of legal proceedings, and the risks detailed in Esperion's filings with the Securities and Exchange Commission. Any forward-looking statements contained in this press release speak only as of the date hereof, and Esperion disclaims any obligation or undertaking to update or revise any forward-looking statements contained in this press release, other than to the extent required by law.

Esperion Contact Information:
Investors:
Alexis Callahan
investorelations@esperion.com
(406) 539-1762

Media:
Tiffany Aldrich
corporateteam@esperion.com
(616) 443-8438